

ELLIPTIC CURVES WITH GOOD REDUCTION AWAY FROM 2: III

RICHARD G.E. PINCH

ABSTRACT. We list the elliptic curves defined over $\mathbb{Q}(\sqrt{5})$ with good reduction away from 2. There are 368 isomorphism classes. We give a global minimal model for each class.

1. INTRODUCTION

In this paper we list the elliptic curves defined over $\mathbb{Q}(\sqrt{5})$ with good reduction away from 2. We use the results of the previous papers [8] and [9], referred to as I and II respectively. By Theorem 1.14 of I, such a curve must have a point of order 2 defined over $\mathbb{Q}(\sqrt{5})$ and by Theorem 2.3 of II, if $t \in \mathbb{Q}(\sqrt{5})$ is the corresponding value of the Hauptmodul on $X_0(2)$ then either t or $t' = 4096/t$ satisfies one of the equations

$$(1.1) \quad t = 64u/v, \quad u + v = x^2$$

or

$$(1.2) \quad t = 64v/2^a u, \quad 2^a u + v = x^2$$

where u, v are units, $x \in \mathbb{Q}(\sqrt{5})$ and $a \geq 0$. We solve these equations and determine the corresponding j -invariants and obtain a global minimal equation in each isomorphism class by Tate's algorithm. There are 368 isomorphism classes of these curves.

2. DIOPHANTINE EQUATIONS OVER THE QUADRATIC FIELD

In this section we solve the Diophantine equations (1.1) and (1.2) over $\mathbb{Q}(\sqrt{5})$. The fundamental unit is $\epsilon = \frac{1+\sqrt{5}}{2}$ and the ring of integers is $\mathbb{Z}[\epsilon]$. Let $\mathcal{N}$ and $\mathcal{T}$ denote absolute norm and trace: let $'$ denote the non-trivial automorphism of $\mathbb{Q}(\sqrt{5})$.

Theorem 2.1. *The equation*

$$2x^2 = u + v, \quad u, v \text{ units of } \mathbb{Q}(\sqrt{5}),$$

has the solutions

a)	$x = 0$	$u = 1$	$v = -1$
b)	1	1	1
c)	1	ϵ^2	ϵ'
d)	3	ϵ^6	ϵ'^6

and all other solutions are obtained by conjugation and scaling by units.

The equations

$$x^2 = 2^a u + v, \quad u, v \text{ units}, \quad a \geq 0,$$

Date: 4 March 1998.

1991 Mathematics Subject Classification. 14K07 (11G05, 14G25).

have the solutions

e)	$a = 0$	$x = 0$	$u = 1$	$v = -1$
f)	0	1	ϵ	ϵ'
g)	0	1	ϵ^2	$-\epsilon$
h)	0	2	ϵ^3	ϵ'^3
i)	1	1	1	-1
j)	1	1	$-\epsilon$	ϵ^3
k)	1	ϵ	1	$-\epsilon'$
l)	1	$\sqrt{5}$	ϵ^2	ϵ'^3
m)	1	$8 + 15\epsilon$	ϵ^{13}	ϵ'
n)	2	$\sqrt{5}$	1	1
o)	2	ϵ^3	ϵ^3	1
p)	3	3	1	1
q)	3	$5 + 2\sqrt{5}$	ϵ^5	1
r)	3	$3 + 2\sqrt{5}$	ϵ^4	1
s)	4	$17 + 8\sqrt{5}$	ϵ^9	1

and all other solutions are obtained by conjugation or scaling by units.

The corresponding values of t are

a,e)	-64
b)	64
c)	$-64\epsilon^3$
d,h)	$-64\epsilon^6$
f)	$-64\epsilon^2$
g)	64ϵ
i)	-32
j)	-32ϵ
k)	$-32\epsilon^2$
l)	$32\epsilon^5$
m)	$-32\epsilon^{14}$
n)	16
o)	$-16\epsilon^3$
p)	8
q)	$-8\epsilon^5$
r)	$8\epsilon^4$
s)	$-4\epsilon^9$

and their conjugates.

Proof. The Theorem will follow from Propositions 2.3, 2.5 and 2.6. $\square$

We define *Lucas numbers* L_n and *Fibonacci numbers* F_n to satisfy the recurrence relation

$$X_{n+1} = X_n + X_{n-1}$$

with the initial conditions $L_0 = 2$, $L_1 = 1$ and $F_0 = 0$, $F_1 = 1$ respectively. Equivalently, we have

$$\epsilon^n = \frac{L_n + F_n\sqrt{5}}{2}$$

We quote the next result from Cohn [2] Theorems 1 and 2.

Proposition 2.2. *The Diophantine equations*

$$\begin{aligned} L_n &= X^2; \\ L_n &= 2X^2 \end{aligned}$$

have the solutions

$$n = 1, 3, \quad X = \pm 1, \pm 3;$$

and

$$n = 0, \pm 6, \quad X = \pm 1, \pm 3$$

respectively.

Proposition 2.3. *The equations*

$$\begin{aligned} 2x^2 &= u + v; \\ x^2 &= u + v \end{aligned}$$

with u, v units, have the solutions

$$\begin{aligned} 2.0^2 &= 1 - 1 \\ 2.1^2 &= 1 + 1 \\ 2.1^2 &= \epsilon^2 + \epsilon' \\ 2.3^2 &= \epsilon^6 + \epsilon'^6 \end{aligned}$$

and

$$\begin{aligned} 0^2 &= 1 - 1 \\ 1^2 &= \epsilon + \epsilon' \\ 1^2 &= \epsilon^2 - \epsilon \\ 2^2 &= \epsilon^3 + \epsilon'^3 \end{aligned}$$

respectively.

Proof. Let A be 1 or 2, according to the equation we consider. Put $U = \mathcal{N}u$, $V = \mathcal{N}v$, $X = \mathcal{N}x$, $w = uv'$. Taking norms we have

$$A^2 X^2 = U + V + \mathcal{T}w$$

If $U = -V$ then $\mathcal{T}w = (AX)^2$, so that by Proposition 2.2 we have $AX = \pm 1$ or ± 2 . But 2 is not a norm, so X must be 1, giving $w = \epsilon$ or ϵ^3 , and solutions $1^2 = \epsilon^2 - \epsilon$ and $2.1^2 = \epsilon^2 + \epsilon'$.

If $U = V$, then $u^{-1} = Uu'$, $v^{-1} = Vv'$. Now

$$Ax'^2 = u' + v' = U(u^{-1} + v^{-1}) = \frac{U}{uv}(u + v) = \frac{U}{uv}Ax^2,$$

so that

$$x'^2 = \frac{U}{uv}x^2,$$

and, considering the factorisation of x^2 into prime elements, we have $x^2 = Yw$ where $Y \in \mathbb{Z}$ and w is a unit. Scaling by w , we may take $x = Z$ or $Z\sqrt{5}$ where $Z \in \mathbb{Z}$. But now $x^2 \in \mathbb{Z}$ and so $u - u' = v' - v$. Thus either $u = \pm 1$ and $v = \pm 1$, or $u = v'$. The first case gives us the solutions $A.0^2 = 1 - 1$ and $2.1^2 = 1 + 1$ and the second gives the equation $Ax^2 = \pm L_n$ for some n . Considering signs, $Ax^2 = L_n$. We cannot have $x^2 = 5Z^2$, for then $\sqrt{5}$ would divide ϵ^n ; so $x^2 = Z^2$ and we have $AZ^2 = L_n$. Applying Proposition 2.2 we have the remaining solutions. $\square$

The next result is established in proposition 9 of [10] using techniques of transcendence theory: the method is further developed in [11].

Proposition 2.4. *The pairs of simultaneous Diophantine equations*

$$\begin{aligned}(X+2)^2 - 10Y^2 &= -1, \\ (X-2)^2 - 2Z^2 &= -1;\end{aligned}$$

and

$$\begin{aligned}X^2 - 2Y^2 &= -1, \\ X^2 - 10Z^2 &= -9\end{aligned}$$

have the solutions

$$(X, Y, Z) = (1, \pm 1, \pm 1) \text{ or } (-5, \pm 1, \pm 5);$$

and

$$(X, Y, Z) = (\pm 1, \pm 1, \pm 1) \text{ or } (\pm 41, \pm 29, \pm 13)$$

respectively.

Proposition 2.5. *The equation*

$$x^2 = 2u + v, \quad u, v \text{ units},$$

has the solutions

$$\begin{aligned}1^2 &= 2 \cdot 1 - 1 \\ 1^2 &= 2(-\epsilon) + \epsilon^3 \\ \epsilon^2 &= 2 \cdot 1 - \epsilon' \\ (\sqrt{5})^2 &= 2\epsilon^2 + \epsilon'^3 \\ (8 + 15\epsilon)^2 &= 2\epsilon^1 3 + \epsilon' .\end{aligned}$$

Proof. Let $U = \mathcal{N}u$, $V = \mathcal{N}v$, $w = uv' = \frac{S+P\sqrt{5}}{2}$, $W = \mathcal{N}w$, $X = \mathcal{N}x$. We have $W = UV$,

$$X^2 = 4U + V + 2S$$

and

$$S^2 - 5P^2 = 4W.$$

Clearly X is odd. If $X = 1$, then x is a unit, and we solve the equation by Proposition 2.3, giving the solutions $1^2 = 2 \cdot 1 - 1$, $1^2 = 2(-\epsilon) + \epsilon^3$ and $\epsilon^2 = 2 \cdot 1 - \epsilon'$. We note that X cannot be 3, which is not a norm from $\mathbb{Q}(\sqrt{5})$.

Assume now that $X \geq 5$, and consider four cases according to the values of U and V .

Suppose that $U = V = +1$. Then $W = +1$ and, substituting, we have

$$\left(\frac{X^2 - 5}{2}\right)^2 - 5P^2 = 4,$$

so

$$(X^2 - 1)(X^2 - 9) = 5(2P)^2.$$

Since X is odd, the two terms on the LHS are congruent to 0 and 8 mod 16, so their highest common factor is 8; since $X \geq 5$, each term is positive. We must have either

$$\begin{aligned}X^2 - 1 &= 10Y^2, \\ X^2 - 9 &= 2Z^2;\end{aligned}$$

or

$$\begin{aligned} X^2 - 1 &= 2Y^2, \\ X^2 - 9 &= 10Z^2. \end{aligned}$$

The first pair have no solution: since 2 is not a quadratic residue of 3, 3 must divide X , but then -1 would be a quadratic residue of 3, a contradiction. The second pair lead to no solution of the original equation; for, from the second equation of the pair, $X^2 \equiv 4 \pmod{5}$, so $X \equiv \pm 2 \pmod{5}$, but this is impossible since X is a norm from $\mathbb{Q}(\sqrt{5})$. Hence there is no solution in the case $U = V = +1$.

Suppose now that $U = +1$, $V = -1$, so that $W = -1$. We have

$$\left(\frac{X^2 - 3}{2}\right)^2 - 5P^2 = -4,$$

so

$$((X + 2)^2 + 1)((X - 2)^2 + 1) = 5(2P)^2.$$

Since X is odd, each term on the LHS is congruent to 2 mod (4) and so their highest common factor is 2; further, each is positive. So

$$\begin{aligned} (X + 2)^2 + 1 &= 10Y^2, \\ (X - 2)^2 + 1 &= 2Z^2. \end{aligned}$$

By Proposition 2.4, the solutions are $(X, Y, Z) = (1, \pm 1, \pm 1)$, dealt with above, and $(5, \pm 1, \pm 5)$, leading to $(\sqrt{5})^2 = 2\epsilon^2 + \epsilon'^3$.

Suppose now that $U = -1$, $V = +1$, so that $W = -1$. Substituting,

$$\left(\frac{X^2 + 3}{2}\right)^2 - 5P^2 = -4$$

so

$$((X + 1)^2 + 4)((X - 1)^2 + 4) = 5(2P)^2.$$

Since X is odd, each term on the LHS is divisible by 4. The highest common factor of the terms divides their difference, which is $4X$, and so is either 4 (if X is prime to 5) or 20 (if $5 \mid X$). In either case, we have

$$\begin{aligned} (X + 1)^2 + 4 &= 5Y^2, \\ (X - 1)^2 + 4 &= Z^2. \end{aligned}$$

The second equation implies that $(X, Z) = (2, \pm 2)$ or $(0, \pm 2)$; but X must be odd, so there is no solution in this case.

Finally, suppose that $U = V = -1$, so that $W = +1$. Substituting,

$$\left(\frac{X^2 + 5}{2}\right)^2 - 5P^2 = 4,$$

so

$$(X^2 + 1)(X^2 + 9) = 5(2P)^2.$$

Since X is odd, each term on the LHS is congruent to 2 mod (4) and so their highest common factor is 2; further, each is positive. We must have either

$$\begin{aligned} X^2 + 1 &= 10Y^2, \\ X^2 + 9 &= 2Z^2 \end{aligned}$$

or

$$\begin{aligned} X^2 + 1 &= 2Y^2, \\ X^2 + 9 &= 10Z^2. \end{aligned}$$

In the first case, $X^2 \equiv 4 \pmod{5}$, so that $X \equiv \pm 2 \pmod{5}$; but X is a norm from $\mathbb{Q}(\sqrt{5})$ and this is impossible. In the second case, by Proposition 2.4, the solutions are $(X, Y, Z) = (\pm 1, \pm 1, \pm 1)$, dealt with above, and $(\pm 41, \pm 29, \pm 13)$ giving the solution $x = 8 + 15\epsilon$, $u = \epsilon^{13}$, $v = \epsilon'$. $\square$

Proposition 2.6. *The equation*

$$x^2 = 2^a u + v, \quad a \geq 2, \quad u, v \text{ units},$$

has solutions

$a = 2$	$x = \sqrt{5}$	$u = 1$	$v = 1$
2	ϵ^3	ϵ^3	1
3	3	1	1
3	$5 + 2\sqrt{5}$	ϵ^5	1
3	$3 + 2\sqrt{5}$	ϵ^4	1
4	$17 + 8\sqrt{5}$	ϵ^9	1.

Proof. We have $v \equiv x^2 \pmod{4}$, so v is a square unit; scaling, we may assume that $v = 1$. Now

$$x^2 - 1 = 2^a u,$$

and, factorising and choosing the sign on x , we have

$$\begin{aligned} x + 1 &= 2^{a-1} u_1, \\ x - 1 &= 2u_2, \end{aligned}$$

where u_1, u_2 are units with $u = u_1 u_2$. Subtracting,

$$2^{a-2} u_1 - u_2 = 1,$$

that is,

$$2^{a-2} = u_1^{-1} + u_2 u_1^{-1}.$$

This is solved by Proposition 2.3, taking the solutions with x a power of 2, and the result follows. $\square$

3. EQUATIONS OF CURVES WITH GOOD REDUCTION AWAY FROM 2

By Corollary 1.14.1 of II, an elliptic curve E defined over $\mathbb{Q}(\sqrt{5})$ with good reduction away from 2 has a point P of order 2 defined over $\mathbb{Q}(\sqrt{5})$. By Proposition 2.1 of I, the corresponding invariant j is given by

$$j(t) = \frac{(t + 16)^3}{t}$$

where $t \in \mathbb{Q}(\sqrt{5})$ is the value of the Hauptmodul for $X_0(2)$ corresponding to the pair (E, P) , and the isogenous curve E' has invariant $j' = j(t')$, with $tt' = 4096$.

The possible values of t are obtained from Theorem 2.1. Table 1 lists values of t corresponding to a curve with good reduction away from 2, together with the corresponding values of j and their factorisations. As in Section 5 of I we can list the minimal equation of a representative of each isomorphism class over $\mathbb{Q}(\sqrt{5})$, and these are given in Table 2. We express the curve in the form

$$Y^2 + a_1 XY + a_3 Y = X^3 + a_2 X^2 + a_4 X + a_6$$

and give the coefficients a_i , the discriminant Δ , the exponent of the conductor 2^f and the reduction type according to Tate's algorithm (see Tate [12], Cremona [4], or [7]). In each table *code* refers to the j invariant.

Theorem 3.1. *There are 56 values of t and 43 values of j corresponding to elliptic curves over $\mathbb{Q}(\sqrt{5})$ with good reduction away from 2. There are 368 isomorphism classes of such curves.*

It is interesting to observe that every curve in Table 2 has additive bad reduction. In addition, we see that there is no curve with good reduction everywhere over $\mathbb{Q}(\sqrt{5})$: see Cremona [3], Ishii [5],[6] and Pinch [7].

t	$code$	j
$-220 + 136\epsilon$	$+\epsilon^{-9}2^2$	34
$-84 - 136\epsilon$	$-\epsilon^9 2^2$	34
$-64 + 40\epsilon$	$+\epsilon^{-5}2^3$	28
$40 - 24\epsilon$	$+\epsilon^{-4}2^3$	31
8	$+2^3$	11
$16 + 24\epsilon$	$+\epsilon^4 2^3$	28
$-24 - 40\epsilon$	$-\epsilon^5 2^3$	31
$-48 + 32\epsilon$	$+\epsilon^{-3}2^4$	14
16	$+2^4$	14
$-16 - 32\epsilon$	$-\epsilon^3 2^4$	14
$-19520 + 12064\epsilon$	$-\epsilon^{-14}2^5$	40
256	$-\epsilon^{-5}2^5$	2
$-64 + 32\epsilon$	$-\epsilon^{-2}2^5$	17
$-32 + 32\epsilon$	$+\epsilon^{-1}2^5$	6
-32	-2^5	5
-32ϵ	$-\epsilon^2 2^5$	13
$-32 - 32\epsilon$	$-\epsilon^2 2^5$	1
96	$+\epsilon^5 2^5$	23
$-7456 - 12064\epsilon$	$-\epsilon^{14}2^5$	41
$-832 + 512\epsilon$	$-\epsilon^{-6}2^6$	39
192	$-\epsilon^{-3}2^6$	32
$-192 + 128\epsilon$	$+\epsilon^{-3}2^6$	26
$-128 + 64\epsilon$	$-\epsilon^{-2}2^6$	8
64	$-\epsilon^{-1}2^6$	4
$-64 + 64\epsilon$	$+\epsilon^{-1}2^6$	31
-64	-2^6	11
64	$+2^6$	35
-64ϵ	$-\epsilon^2 2^6$	28
64	$+\epsilon^2 2^6$	27
$-64 - 64\epsilon$	$-\epsilon^2 2^6$	30
$-64 - 128\epsilon$	$-\epsilon^3 2^6$	12
64	$+\epsilon^3 2^6$	16
$-320 - 512\epsilon$	$-\epsilon^6 2^6$	22
$-78080 + 48256\epsilon$	$-\epsilon^{-14}2^7$	42
$-1024 + 640\epsilon$	$+\epsilon^{-5}2^7$	7
$-256 + 128\epsilon$	$-\epsilon^{-2}2^7$	33
128	$-\epsilon^{-1}2^7$	24
-128	-2^7	3
128	$+\epsilon^2 2^7$	10
$-128 - 128\epsilon$	$-\epsilon^2 2^7$	18
$-384 - 640\epsilon$	$-\epsilon^5 2^7$	29
$-29824 - 48256\epsilon$	$-\epsilon^{14}2^7$	43
768	$-\epsilon^{-3}2^8$	36
256	$+2^8$	34
$256 + 512\epsilon$	$+\epsilon^3 2^8$	20
4096	$-\epsilon^{-5}2^9$	15
$2560 - 1536\epsilon$	$+\epsilon^{-4}2^9$	38
512	$+2^9$	19
$1024 + 1536\epsilon$	$+\epsilon^4 2^9$	21
$1536 + 2560\epsilon$	$+\epsilon^5 2^9$	37
$56320 - 34816\epsilon$	$-\epsilon^{-9}2^{10}$	25
21504	$+\epsilon^9 2^{10}$	9
78608	$+2^4(17)^3$	
78608	$+2^4(17)^3$	
$4928 + 960\epsilon$	$+\epsilon^2 2^6(5 - \epsilon)^3$	
$5888 - 960\epsilon$	$+\epsilon^{-2}2^6(4 + \epsilon)^3$	
1728	$+2^6(3)^3$	
$4928 + 960\epsilon$	$+\epsilon^2 2^6(5 - \epsilon)^3$	
$5888 - 960\epsilon$	$+\epsilon^{-2}2^6(4 + \epsilon)^3$	
2048	$+2^{11}$	
2048	$+2^{11}$	
2048	$+2^{11}$	
$525604480 - 324907648\epsilon$	$-\epsilon^{-7}2^7(50 - 13\epsilon)^3$	
$103808 - 64640\epsilon$	$-\epsilon^{-7}2^7(4 + \epsilon)^3$	
$2688 - 1664\epsilon$	$-\epsilon^{-7}2^7$	
$1280 + 640\epsilon$	$-\epsilon^2 2^7(1 - 2\epsilon)^3$	
128	$+2^7$	
$1920 - 640\epsilon$	$-\epsilon^{-1}2^7(1 - 2\epsilon)^3$	
$1024 + 1664\epsilon$	$+\epsilon^7 2^7$	
$39168 + 64640\epsilon$	$+\epsilon^7 2^7(5 - \epsilon)^3$	
$200696832 + 324907648\epsilon$	$+\epsilon^7 2^7(37 + 13\epsilon)^3$	
$914880 - 565760\epsilon$	$+\epsilon^{-12}2^6(-1 - 8\epsilon)^3$	
$63168 - 39040\epsilon$	$-\epsilon^{-15}2^6$	
$44864 - 26496\epsilon$	$+\epsilon^{-6}2^6(5 + 2\epsilon)^3$	
$15040 - 9280\epsilon$	$-\epsilon^{-7}2^6(1 - 2\epsilon)^3$	
$12032 - 7232\epsilon$	$+\epsilon^{-8}2^6(3 + 2\epsilon)^3$	
$5888 - 960\epsilon$	$+\epsilon^{-2}2^6(4 + \epsilon)^3$	
1728	$+2^6(3)^3$	
8000	$+2^6(5)^3$	
$4928 + 960\epsilon$	$+\epsilon^2 2^6(5 - \epsilon)^3$	
$4800 + 7232\epsilon$	$+\epsilon^8 2^6(5 - 2\epsilon)^3$	
$5760 + 9280\epsilon$	$-\epsilon^7 2^6(1 - 2\epsilon)^3$	
$18368 + 26496\epsilon$	$+\epsilon^6 2^6(7 - 2\epsilon)^3$	
$24128 + 39040\epsilon$	$+\epsilon^{15}2^6$	
$349120 + 565760\epsilon$	$-\epsilon^{12}2^6(9 - 8\epsilon)^3$	
$8421373408 - 5204711200\epsilon$	$+\epsilon^{-13}2^5(-85 + 6\epsilon)^3$	
$1409888 - 870240\epsilon$	$+\epsilon^{-4}2^5(17 - 6\epsilon)^3$	
$70368 - 43040\epsilon$	$+\epsilon^{-4}2^5(7 - \epsilon)^3$	
$39680 - 22560\epsilon$	$-\epsilon^2 2^5(9 - 8\epsilon)^3$	
10976	$+2^5(7)^3$	
$17120 + 22560\epsilon$	$-\epsilon^{-1}2^5(-1 - 8\epsilon)^3$	
$27328 + 43040\epsilon$	$+\epsilon^4 2^5(6 + \epsilon)^3$	
$539648 + 870240\epsilon$	$+\epsilon^4 2^5(11 + 6\epsilon)^3$	
$3216662208 + 5204711200\epsilon$	$-\epsilon^{13}2^5(-79 - 6\epsilon)^3$	
$889584 - 548896\epsilon$	$+\epsilon^{-6}2^4(15 - 2\epsilon)^3$	
78608	$+2^4(17)^3$	
$340688 + 548896\epsilon$	$+\epsilon^6 2^4(13 + 2\epsilon)^3$	
$23528168 - 14540840\epsilon$	$-\epsilon^{-10}2^3(-29 + 5\epsilon)^3$	
$9036560 - 5578728\epsilon$	$-\epsilon^{-8}2^3(-34 - 3\epsilon)^3$	
287496	$+2^3(33)^3$	
$3457832 + 5578728\epsilon$	$-\epsilon^8 2^3(-37 + 3\epsilon)^3$	
$8987328 + 14540840\epsilon$	$-\epsilon^{10}2^3(-24 - 5\epsilon)^3$	
$4386800300 - 2711191688\epsilon$	$-\epsilon^{-21}2^2(34 - 9\epsilon)^3$	
$1675608612 + 2711191688\epsilon$	$+\epsilon^{21}2^2(25 + 9\epsilon)^3$	

Table 1. Invariants of elliptic curves defined over $\mathbb{Q}(\sqrt{5})$ with good reduction away from 2.

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$-\epsilon$	0	$-16 + 11\epsilon$	$27 - 17\epsilon$	$-\epsilon^{-3}2^8$	I_1^*	3	36
0	$-\epsilon$	0	$-171 + 106\epsilon$	$-1050 + 647\epsilon$	$-\epsilon^3 2^{10}$	III^*	3	25
0	$-1 + \epsilon$	0	$-\epsilon$	0	$+\epsilon^6 2^4$	III	3	14
0	$-1 + \epsilon$	0	$-5 - 6\epsilon$	$-7 - 11\epsilon$	$+\epsilon^6 2^8$	I_1^*	3	34
0	$-1 + \epsilon$	0	$-5 - 11\epsilon$	$10 + 17\epsilon$	$+\epsilon^3 2^8$	I_3^*	3	20
0	$-1 + \epsilon$	0	$-65 - 106\epsilon$	$-403 - 647\epsilon$	$+\epsilon^{-3} 2^{10}$	III^*	3	9
0	ϵ	0	$-16 + 11\epsilon$	$-27 + 17\epsilon$	$-\epsilon^{-3} 2^8$	I_0^*	4	36
0	ϵ	0	$-171 + 106\epsilon$	$1050 - 647\epsilon$	$-\epsilon^3 2^{10}$	I_2^*	4	25
0	1	0	$-171 - 277\epsilon$	$-1697 - 2747\epsilon$	$+\epsilon^3 2^{10}$	I_2^*	4	9
0	1	0	$-448 + 277\epsilon$	$-4444 + 2747\epsilon$	$-\epsilon^{-3} 2^{10}$	I_2^*	4	25
0	$1 + \epsilon$	0	ϵ	0	$+2^4$	II	4	14
0	$1 + \epsilon$	0	-4ϵ	$-4 - 8\epsilon$	$+\epsilon^{-3} 2^8$	I_0^*	4	20
0	$1 + \epsilon$	0	$-5 + \epsilon$	$-4 + 3\epsilon$	$+2^8$	I_0^*	4	34
0	$1 + \epsilon$	0	$-5 + 6\epsilon$	$7 - 3\epsilon$	$-\epsilon^3 2^8$	I_0^*	4	36
0	$1 - \epsilon$	0	$-\epsilon$	0	$+\epsilon^6 2^4$	II	4	14
0	$1 - \epsilon$	0	$-5 - 6\epsilon$	$7 + 11\epsilon$	$+\epsilon^6 2^8$	I_0^*	4	34
0	$1 - \epsilon$	0	$-5 - 11\epsilon$	$-10 - 17\epsilon$	$+\epsilon^3 2^8$	I_0^*	4	20
0	$1 - \epsilon$	0	$-65 - 106\epsilon$	$403 + 647\epsilon$	$+\epsilon^{-3} 2^{10}$	I_2^*	4	9
0	-1	0	$-171 - 277\epsilon$	$1697 + 2747\epsilon$	$+\epsilon^3 2^{10}$	I_2^*	4	9
0	-1	0	$-448 + 277\epsilon$	$4444 - 2747\epsilon$	$-\epsilon^{-3} 2^{10}$	I_2^*	4	25
0	$-1 - \epsilon$	0	ϵ	0	$+2^4$	II	4	14
0	$-1 - \epsilon$	0	-4ϵ	$4 + 8\epsilon$	$+\epsilon^{-3} 2^8$	I_0^*	4	20
0	$-1 - \epsilon$	0	$-5 + \epsilon$	$4 - 3\epsilon$	$+2^8$	I_0^*	4	34
0	$-1 - \epsilon$	0	$-5 + 6\epsilon$	$-7 + 3\epsilon$	$-\epsilon^3 2^8$	I_0^*	4	36
0	0	0	4	0	-2^{12}	I_3^*	5	11
0	0	0	$4 + 4\epsilon$	0	$-\epsilon^6 2^{12}$	I_3^*	5	11
0	0	0	-1	0	$+2^6$	III	5	11
0	0	0	$-1 - \epsilon$	0	$+\epsilon^6 2^6$	III	5	11
0	0	0	-11	14	$+2^9$	I_0^*	5	19
0	0	0	-11	-14	$+2^9$	I_0^*	5	19
0	0	0	$-11 - 11\epsilon$	$14 + 28\epsilon$	$+\epsilon^6 2^9$	I_0^*	5	19
0	0	0	$-11 - 11\epsilon$	$-14 - 28\epsilon$	$+\epsilon^6 2^9$	I_0^*	5	19
0	ϵ	0	-1	$-\epsilon$	$+\epsilon^2 2^6$	III	5	31
0	ϵ	0	$-1 - 5\epsilon$	$3 + 2\epsilon$	$+\epsilon^2 2^{12}$	I_3^*	5	27
0	ϵ	0	$-11 - 10\epsilon$	$-22 - 31\epsilon$	$+\epsilon^{-2} 2^9$	I_0^*	5	21
0	ϵ	0	$-32 - 53\epsilon$	$-155 - 251\epsilon$	$+\epsilon^5 2^9$	I_0^*	5	37
0	$-\epsilon$	0	-1	ϵ	$+\epsilon^2 2^6$	III	5	31
0	$-\epsilon$	0	$-1 - 5\epsilon$	$-3 - 2\epsilon$	$+\epsilon^2 2^{12}$	I_3^*	5	27
0	$-\epsilon$	0	$-11 - 10\epsilon$	$22 + 31\epsilon$	$+\epsilon^{-2} 2^9$	I_0^*	5	21
0	$-\epsilon$	0	$-32 - 53\epsilon$	$155 + 251\epsilon$	$+\epsilon^5 2^9$	I_0^*	5	37
0	1	0	$3 - 4\epsilon$	$-5 + 4\epsilon$	$+\epsilon^{-5} 2^{12}$	I_3^*	5	27
0	1	0	$-1 + 4\epsilon$	$-1 - 4\epsilon$	$-\epsilon^5 2^{12}$	I_3^*	5	4
0	1	0	$-1 - \epsilon$	$-1 - \epsilon$	$+\epsilon^4 2^6$	III	5	28
0	1	0	$-2 + \epsilon$	$-2 + \epsilon$	$+\epsilon^{-4} 2^6$	III	5	31
0	1	0	$-11 - 21\epsilon$	$-37 - 59\epsilon$	$+\epsilon^{-1} 2^9$	I_0^*	5	37
0	1	0	$-32 + 21\epsilon$	$-96 + 59\epsilon$	$-\epsilon^2 2^9$	I_0^*	5	15
0	$1 + \epsilon$	0	$-21 - 31\epsilon$	$-84 - 137\epsilon$	$+\epsilon^4 2^9$	I_0^*	5	21
0	$1 + \epsilon$	0	$-53 + 33\epsilon$	$168 - 105\epsilon$	$+\epsilon^{-4} 2^9$	I_0^*	5	38

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$1 - \epsilon$	0	-1	$-1 + \epsilon$	$+\epsilon^{-2}2^6$	III	5	28
0	$1 - \epsilon$	0	$-6 + 5\epsilon$	$5 - 2\epsilon$	$-\epsilon^{-1}2^{12}$	I_3^*	5	4
0	$1 - \epsilon$	0	$-21 + 10\epsilon$	$-53 + 31\epsilon$	$+\epsilon^22^9$	I_0^*	5	38
0	$1 - \epsilon$	0	$-85 + 53\epsilon$	$-406 + 251\epsilon$	$-\epsilon^{-5}2^9$	I_0^*	5	15
0	-1	0	$3 - 4\epsilon$	$5 - 4\epsilon$	$+\epsilon^{-5}2^{12}$	I_3^*	5	27
0	-1	0	$-1 + 4\epsilon$	$1 + 4\epsilon$	$-\epsilon^52^{12}$	I_3^*	5	4
0	-1	0	$-1 - \epsilon$	$1 + \epsilon$	$+\epsilon^42^6$	III	5	28
0	-1	0	$-2 + \epsilon$	$2 - \epsilon$	$+\epsilon^{-4}2^6$	III	5	31
0	-1	0	$-11 - 21\epsilon$	$37 + 59\epsilon$	$+\epsilon^{-1}2^9$	I_0^*	5	37
0	-1	0	$-32 + 21\epsilon$	$96 - 59\epsilon$	$-\epsilon^22^9$	I_0^*	5	15
0	$-1 + \epsilon$	0	-1	$1 - \epsilon$	$+\epsilon^{-2}2^6$	III	5	28
0	$-1 + \epsilon$	0	$-6 + 5\epsilon$	$-5 + 2\epsilon$	$-\epsilon^{-1}2^{12}$	I_3^*	5	4
0	$-1 + \epsilon$	0	$-21 + 10\epsilon$	$53 - 31\epsilon$	$+\epsilon^22^9$	I_0^*	5	38
0	$-1 + \epsilon$	0	$-85 + 53\epsilon$	$406 - 251\epsilon$	$-\epsilon^{-5}2^9$	I_0^*	5	15
0	$-1 - \epsilon$	0	$-21 - 31\epsilon$	$84 + 137\epsilon$	$+\epsilon^42^9$	I_0^*	5	21
0	$-1 - \epsilon$	0	$-53 + 33\epsilon$	$-168 + 105\epsilon$	$+\epsilon^{-4}2^9$	I_0^*	5	38
0	0	0	ϵ	0	$-\epsilon^32^6$	II	6	11
0	0	0	4ϵ	0	$-\epsilon^32^{12}$	I_2^*	6	11
0	0	0	$-\epsilon$	0	$+\epsilon^32^6$	II	6	11
0	0	0	-4ϵ	0	$+\epsilon^32^{12}$	I_2^*	6	11
0	0	0	1	0	-2^6	II	6	11
0	0	0	$1 + \epsilon$	0	$-\epsilon^62^6$	II	6	11
0	0	0	$1 - \epsilon$	0	$+\epsilon^{-3}2^6$	II	6	11
0	0	0	$4 - 4\epsilon$	0	$+\epsilon^{-3}2^{12}$	I_2^*	6	11
0	0	0	$-1 + \epsilon$	0	$-\epsilon^{-3}2^6$	II	6	11
0	0	0	-4	0	$+2^{12}$	I_2^*	6	11
0	0	0	$-4 + 4\epsilon$	0	$-\epsilon^{-3}2^{12}$	I_2^*	6	11
0	0	0	$-4 - 4\epsilon$	0	$+\epsilon^62^{12}$	I_2^*	6	11
0	0	0	-44	112	$+2^{15}$	I_5^*	6	19
0	0	0	-44	-112	$+2^{15}$	I_5^*	6	19
0	0	0	$-44 - 44\epsilon$	$112 + 224\epsilon$	$+\epsilon^62^{15}$	I_5^*	6	19
0	0	0	$-44 - 44\epsilon$	$-112 - 224\epsilon$	$+\epsilon^62^{15}$	I_5^*	6	19
0	ϵ	0	$-\epsilon$	$-1 - \epsilon$	$+\epsilon^22^6$	II	6	27
0	ϵ	0	$-1 - 2\epsilon$	$-2 - 3\epsilon$	$+\epsilon^52^6$	II	6	30
0	ϵ	0	$-4 - 6\epsilon$	$4 + 6\epsilon$	$+\epsilon^32^6$	II	6	22
0	ϵ	0	$-5 - \epsilon$	$-1 + 2\epsilon$	$+\epsilon^22^{12}$	I_2^*	6	31
0	ϵ	0	$-5 - 9\epsilon$	$7 + 10\epsilon$	$+\epsilon^52^{12}$	I_2^*	6	30
0	ϵ	0	$-17 - 25\epsilon$	$-57 - 90\epsilon$	$+\epsilon^32^{12}$	I_2^*	6	22
0	ϵ	0	$-45 - 41\epsilon$	$135 + 162\epsilon$	$+\epsilon^{-2}2^{15}$	I_5^*	6	21
0	ϵ	0	$-65 + 43\epsilon$	$259 - 158\epsilon$	$-\epsilon^{-3}2^{14}$	I_4^*	6	36
0	ϵ	0	$-129 - 213\epsilon$	$1027 + 1666\epsilon$	$+\epsilon^52^{15}$	I_5^*	6	37
0	ϵ	0	$-685 + 423\epsilon$	$-7977 + 4914\epsilon$	$-\epsilon^32^{16}$	I_6^*	6	25
0	$-\epsilon$	0	$-\epsilon$	$1 + \epsilon$	$+\epsilon^22^6$	II	6	27
0	$-\epsilon$	0	$-1 - 2\epsilon$	$2 + 3\epsilon$	$+\epsilon^52^6$	II	6	30
0	$-\epsilon$	0	$-4 - 6\epsilon$	$-4 - 6\epsilon$	$+\epsilon^32^6$	II	6	22
0	$-\epsilon$	0	$-5 - \epsilon$	$1 - 2\epsilon$	$+\epsilon^22^{12}$	I_2^*	6	31
0	$-\epsilon$	0	$-5 - 9\epsilon$	$-7 - 10\epsilon$	$+\epsilon^52^{12}$	I_2^*	6	30
0	$-\epsilon$	0	$-17 - 25\epsilon$	$57 + 90\epsilon$	$+\epsilon^32^{12}$	I_2^*	6	22

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$-\epsilon$	0	$-45 - 41\epsilon$	$-135 - 162\epsilon$	$+\epsilon^{-2}2^{15}$	I_5^*	6	21
0	$-\epsilon$	0	$-65 + 43\epsilon$	$-259 + 158\epsilon$	$-\epsilon^{-3}2^{14}$	I_4^*	6	36
0	$-\epsilon$	0	$-129 - 213\epsilon$	$-1027 - 1666\epsilon$	$+\epsilon^5 2^{15}$	I_5^*	6	37
0	$-\epsilon$	0	$-685 + 423\epsilon$	$7977 - 4914\epsilon$	$-\epsilon^3 2^{16}$	I_6^*	6	25
0	1	0	ϵ	ϵ	$-\epsilon^5 2^6$	II	6	4
0	1	0	$-\epsilon$	$-\epsilon$	$+\epsilon^{-1}2^6$	II	6	30
0	1	0	$1 - \epsilon$	$1 - \epsilon$	$+\epsilon^{-5}2^6$	II	6	27
0	1	0	$-1 + \epsilon$	$-1 + \epsilon$	$-\epsilon 2^6$	II	6	8
0	1	0	$-1 - 4\epsilon$	$-1 + 4\epsilon$	$+\epsilon^{-1}2^{12}$	I_2^*	6	30
0	1	0	$-2 - 2\epsilon$	2ϵ	$+\epsilon^{-3}2^6$	II	6	22
0	1	0	$-4 + 2\epsilon$	$2 - 2\epsilon$	$-\epsilon^3 2^6$	II	6	39
0	1	0	$-5 + 4\epsilon$	$3 - 4\epsilon$	$-\epsilon 2^{12}$	I_2^*	6	8
0	1	0	$-5 - 4\epsilon$	$3 + 4\epsilon$	$+\epsilon^4 2^{12}$	I_2^*	6	28
0	1	0	$-9 + 4\epsilon$	$7 - 4\epsilon$	$+\epsilon^{-4}2^{12}$	I_2^*	6	31
0	1	0	$-9 - 8\epsilon$	$-9 - 24\epsilon$	$+\epsilon^{-3}2^{12}$	I_2^*	6	22
0	1	0	$-17 + 8\epsilon$	$-33 + 24\epsilon$	$-\epsilon^3 2^{12}$	I_2^*	6	39
0	1	0	$-45 - 84\epsilon$	$251 + 388\epsilon$	$+\epsilon^{-1}2^{15}$	I_5^*	6	37
0	1	0	$-129 + 84\epsilon$	$639 - 388\epsilon$	$-\epsilon 2^{15}$	I_5^*	6	15
0	1	0	$-685 - 1108\epsilon$	$12891 + 20868\epsilon$	$+\epsilon^3 2^{16}$	I_6^*	6	9
0	1	0	$-1793 + 1108\epsilon$	$33759 - 20868\epsilon$	$-\epsilon^{-3}2^{16}$	I_6^*	6	25
0	$1 + \epsilon$	0	$-2 + \epsilon$	-1	$+2^{10}$	I_0^*	6	14
0	$1 + \epsilon$	0	$-2 - 19\epsilon$	$11 + 24\epsilon$	$+\epsilon^{-3}2^{14}$	I_4^*	6	20
0	$1 + \epsilon$	0	$-22 + \epsilon$	$11 - 44\epsilon$	$+2^{14}$	I_4^*	6	34
0	$1 + \epsilon$	0	$-22 + 21\epsilon$	$-57 + 44\epsilon$	$-\epsilon^3 2^{14}$	I_4^*	6	36
0	$1 + \epsilon$	0	$-86 - 127\epsilon$	$459 + 756\epsilon$	$+\epsilon^4 2^{15}$	I_5^*	6	21
0	$1 + \epsilon$	0	$-214 + 129\epsilon$	$-1429 + 884\epsilon$	$+\epsilon^{-4}2^{15}$	I_5^*	6	38
0	$1 - \epsilon$	0	$-1 + \epsilon$	$-2 + \epsilon$	$-\epsilon^{-1}2^6$	II	6	4
0	$1 - \epsilon$	0	$-2 - 3\epsilon$	$1 + 2\epsilon$	$+\epsilon^6 2^{10}$	I_0^*	6	14
0	$1 - \epsilon$	0	$-3 + 2\epsilon$	$-5 + 3\epsilon$	$-\epsilon^{-5}2^6$	II	6	8
0	$1 - \epsilon$	0	$-6 + \epsilon$	$1 - 2\epsilon$	$+\epsilon^{-2}2^{12}$	I_2^*	6	28
0	$1 - \epsilon$	0	$-10 + 6\epsilon$	$10 - 6\epsilon$	$-\epsilon^{-3}2^6$	II	6	39
0	$1 - \epsilon$	0	$-14 + 9\epsilon$	$17 - 10\epsilon$	$-\epsilon^{-5}2^{12}$	I_2^*	6	8
0	$1 - \epsilon$	0	$-22 - 23\epsilon$	$-55 - 66\epsilon$	$+\epsilon^6 2^{14}$	I_4^*	6	34
0	$1 - \epsilon$	0	$-22 - 43\epsilon$	$101 + 158\epsilon$	$+\epsilon^3 2^{14}$	I_4^*	6	20
0	$1 - \epsilon$	0	$-42 + 25\epsilon$	$-147 + 90\epsilon$	$-\epsilon^{-3}2^{12}$	I_2^*	6	39
0	$1 - \epsilon$	0	$-86 + 41\epsilon$	$297 - 162\epsilon$	$+\epsilon^2 2^{15}$	I_5^*	6	38
0	$1 - \epsilon$	0	$-262 - 423\epsilon$	$-3063 - 4914\epsilon$	$+\epsilon^{-3}2^{16}$	I_6^*	6	9
0	$1 - \epsilon$	0	$-342 + 213\epsilon$	$2693 - 1666\epsilon$	$-\epsilon^{-5}2^{15}$	I_5^*	6	15
0	-1	0	ϵ	$-\epsilon$	$-\epsilon^5 2^6$	II	6	4
0	-1	0	$-\epsilon$	ϵ	$+\epsilon^{-1}2^6$	II	6	30
0	-1	0	$1 - \epsilon$	$-1 + \epsilon$	$+\epsilon^{-5}2^6$	II	6	27
0	-1	0	$-1 + \epsilon$	$1 - \epsilon$	$-\epsilon 2^6$	II	6	8
0	-1	0	$-1 - 4\epsilon$	$1 - 4\epsilon$	$+\epsilon^{-1}2^{12}$	I_2^*	6	30
0	-1	0	$-2 - 2\epsilon$	-2ϵ	$+\epsilon^{-3}2^6$	II	6	22
0	-1	0	$-4 + 2\epsilon$	$-2 + 2\epsilon$	$-\epsilon^3 2^6$	II	6	39
0	-1	0	$-5 + 4\epsilon$	$-3 + 4\epsilon$	$-\epsilon 2^{12}$	I_2^*	6	8
0	-1	0	$-5 - 4\epsilon$	$-3 - 4\epsilon$	$+\epsilon^4 2^{12}$	I_2^*	6	28
0	-1	0	$-9 + 4\epsilon$	$-7 + 4\epsilon$	$+\epsilon^{-4}2^{12}$	I_2^*	6	31

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	-1	0	$-9 - 8\epsilon$	$9 + 24\epsilon$	$+\epsilon^{-3}2^{12}$	I_2^*	6	22
0	-1	0	$-17 + 8\epsilon$	$33 - 24\epsilon$	$-\epsilon^3 2^{12}$	I_2^*	6	39
0	-1	0	$-45 - 84\epsilon$	$-251 - 388\epsilon$	$+\epsilon^{-1}2^{15}$	I_5^*	6	37
0	-1	0	$-129 + 84\epsilon$	$-639 + 388\epsilon$	$-\epsilon 2^{15}$	I_5^*	6	15
0	-1	0	$-685 - 1108\epsilon$	$-12891 - 20868\epsilon$	$+\epsilon^3 2^{16}$	I_6^*	6	9
0	-1	0	$-1793 + 1108\epsilon$	$-33759 + 20868\epsilon$	$-\epsilon^{-3}2^{16}$	I_6^*	6	25
0	$-1 + \epsilon$	0	$-1 + \epsilon$	$2 - \epsilon$	$-\epsilon^{-1}2^6$	II	6	4
0	$-1 + \epsilon$	0	$-2 - 3\epsilon$	$-1 - 2\epsilon$	$+\epsilon^6 2^{10}$	I_0^*	6	14
0	$-1 + \epsilon$	0	$-3 + 2\epsilon$	$5 - 3\epsilon$	$-\epsilon^{-5}2^6$	II	6	8
0	$-1 + \epsilon$	0	$-6 + \epsilon$	$-1 + 2\epsilon$	$+\epsilon^{-2}2^{12}$	I_2^*	6	28
0	$-1 + \epsilon$	0	$-10 + 6\epsilon$	$-10 + 6\epsilon$	$-\epsilon^{-3}2^6$	II	6	39
0	$-1 + \epsilon$	0	$-14 + 9\epsilon$	$-17 + 10\epsilon$	$-\epsilon^{-5}2^{12}$	I_2^*	6	8
0	$-1 + \epsilon$	0	$-22 - 23\epsilon$	$55 + 66\epsilon$	$+\epsilon^6 2^{14}$	I_4^*	6	34
0	$-1 + \epsilon$	0	$-22 - 43\epsilon$	$-101 - 158\epsilon$	$+\epsilon^3 2^{14}$	I_4^*	6	20
0	$-1 + \epsilon$	0	$-42 + 25\epsilon$	$147 - 90\epsilon$	$-\epsilon^{-3}2^{12}$	I_2^*	6	39
0	$-1 + \epsilon$	0	$-86 + 41\epsilon$	$-297 + 162\epsilon$	$+\epsilon^2 2^{15}$	I_5^*	6	38
0	$-1 + \epsilon$	0	$-262 - 423\epsilon$	$3063 + 4914\epsilon$	$+\epsilon^{-3}2^{16}$	I_6^*	6	9
0	$-1 + \epsilon$	0	$-342 + 213\epsilon$	$-2693 + 1666\epsilon$	$-\epsilon^{-5}2^{15}$	I_5^*	6	15
0	$-1 - \epsilon$	0	$-2 + \epsilon$	1	$+2^{10}$	I_0^*	6	14
0	$-1 - \epsilon$	0	$-2 - 19\epsilon$	$-11 - 24\epsilon$	$+\epsilon^{-3}2^{14}$	I_4^*	6	20
0	$-1 - \epsilon$	0	$-22 + \epsilon$	$-11 + 44\epsilon$	$+2^{14}$	I_4^*	6	34
0	$-1 - \epsilon$	0	$-22 + 21\epsilon$	$57 - 44\epsilon$	$-\epsilon^3 2^{14}$	I_4^*	6	36
0	$-1 - \epsilon$	0	$-86 - 127\epsilon$	$-459 - 756\epsilon$	$+\epsilon^4 2^{15}$	I_5^*	6	21
0	$-1 - \epsilon$	0	$-214 + 129\epsilon$	$1429 - 884\epsilon$	$+\epsilon^{-4}2^{15}$	I_5^*	6	38
0	ϵ	0	-2ϵ	$-2 - 2\epsilon$	$+\epsilon^{-1}2^7$	II	7	18
0	ϵ	0	$1 + \epsilon$	$1 + 2\epsilon$	$-\epsilon^6 2^8$	III	7	5
0	ϵ	0	$1 - \epsilon$	-1	$+\epsilon^{-1}2^8$	III	7	6
0	ϵ	0	$3 + 3\epsilon$	$-5 - 10\epsilon$	$-\epsilon^6 2^{14}$	III^*	7	5
0	ϵ	0	$3 - 5\epsilon$	$3 - 2\epsilon$	$+\epsilon^{-1}2^{14}$	III^*	7	6
0	ϵ	0	$-1 + \epsilon$	1	$-\epsilon 2^8$	III	7	17
0	ϵ	0	$-1 - 9\epsilon$	$7 + 6\epsilon$	$+\epsilon^{-1}2^{13}$	I_2^*	7	18
0	ϵ	0	-2	-2ϵ	$+\epsilon^{-2}2^7$	II	7	10
0	ϵ	0	$-2 - 2\epsilon$	$-2 - 4\epsilon$	$+\epsilon^6 2^7$	II	7	3
0	ϵ	0	$-5 + 3\epsilon$	$-5 - 2\epsilon$	$-\epsilon 2^{14}$	III^*	7	17
0	ϵ	0	$-9 - \epsilon$	$-1 + 6\epsilon$	$+\epsilon^{-2}2^{13}$	I_2^*	7	10
0	ϵ	0	$-9 - 9\epsilon$	$7 + 14\epsilon$	$+\epsilon^6 2^{13}$	I_2^*	7	3
0	ϵ	0	$-11 + 7\epsilon$	$19 - 12\epsilon$	$-\epsilon^{-5}2^8$	III	7	2
0	ϵ	0	$-13 + 8\epsilon$	$24 - 15\epsilon$	$+\epsilon^{-2}2^7$	II	7	7
0	ϵ	0	$-45 + 27\epsilon$	$-125 + 78\epsilon$	$-\epsilon^{-5}2^{14}$	III^*	7	2
0	ϵ	0	$-53 + 31\epsilon$	$-161 + 98\epsilon$	$+\epsilon^{-2}2^{13}$	I_2^*	7	7
0	ϵ	0	$-75 - 117\epsilon$	$419 + 680\epsilon$	$+\epsilon^5 2^8$	III	7	41
0	ϵ	0	$-301 - 469\epsilon$	$-3821 - 6210\epsilon$	$+\epsilon^5 2^{14}$	III^*	7	41
0	ϵ	0	$-384 + 237\epsilon$	$-3325 + 2055\epsilon$	$-\epsilon^{-5}2^7$	II	7	42
0	ϵ	0	$-1537 + 947\epsilon$	$27547 - 17030\epsilon$	$-\epsilon^{-5}2^{13}$	I_2^*	7	42
0	$-\epsilon$	0	-2ϵ	$2 + 2\epsilon$	$+\epsilon^{-1}2^7$	II	7	18
0	$-\epsilon$	0	$1 + \epsilon$	$-1 - 2\epsilon$	$-\epsilon^6 2^8$	III	7	5
0	$-\epsilon$	0	$1 - \epsilon$	1	$+\epsilon^{-1}2^8$	III	7	6
0	$-\epsilon$	0	$3 + 3\epsilon$	$5 + 10\epsilon$	$-\epsilon^6 2^{14}$	III^*	7	5

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$-\epsilon$	0	$3 - 5\epsilon$	$-3 + 2\epsilon$	$+\epsilon^{-1}2^{14}$	III^*	7	6
0	$-\epsilon$	0	$-1 + \epsilon$	-1	$-\epsilon 2^8$	III	7	17
0	$-\epsilon$	0	$-1 - 9\epsilon$	$-7 - 6\epsilon$	$+\epsilon^{-1}2^{13}$	I_2^*	7	18
0	$-\epsilon$	0	-2	2ϵ	$+\epsilon^{-2}2^7$	II	7	10
0	$-\epsilon$	0	$-2 - 2\epsilon$	$2 + 4\epsilon$	$+\epsilon^6 2^7$	II	7	3
0	$-\epsilon$	0	$-5 + 3\epsilon$	$5 + 2\epsilon$	$-\epsilon 2^{14}$	III^*	7	17
0	$-\epsilon$	0	$-9 - \epsilon$	$1 - 6\epsilon$	$+\epsilon^{-2}2^{13}$	I_2^*	7	10
0	$-\epsilon$	0	$-9 - 9\epsilon$	$-7 - 14\epsilon$	$+\epsilon^6 2^{13}$	I_2^*	7	3
0	$-\epsilon$	0	$-11 + 7\epsilon$	$-19 + 12\epsilon$	$-\epsilon^{-5}2^8$	III	7	2
0	$-\epsilon$	0	$-13 + 8\epsilon$	$-24 + 15\epsilon$	$+\epsilon^{-2}2^7$	II	7	7
0	$-\epsilon$	0	$-45 + 27\epsilon$	$125 - 78\epsilon$	$-\epsilon^{-5}2^{14}$	III^*	7	2
0	$-\epsilon$	0	$-53 + 31\epsilon$	$161 - 98\epsilon$	$+\epsilon^{-2}2^{13}$	I_2^*	7	7
0	$-\epsilon$	0	$-75 - 117\epsilon$	$-419 - 680\epsilon$	$+\epsilon^5 2^8$	III	7	41
0	$-\epsilon$	0	$-301 - 469\epsilon$	$3821 + 6210\epsilon$	$+\epsilon^5 2^{14}$	III^*	7	41
0	$-\epsilon$	0	$-384 + 237\epsilon$	$3325 - 2055\epsilon$	$-\epsilon^{-5}2^7$	II	7	42
0	$-\epsilon$	0	$-1537 + 947\epsilon$	$-27547 + 17030\epsilon$	$-\epsilon^{-5}2^{13}$	I_2^*	7	42
0	1	0	1	1	-2^8	III	7	5
0	1	0	3	-5	-2^{14}	III^*	7	5
0	1	0	$-1 - 2\epsilon$	$-1 - 2\epsilon$	$+\epsilon^5 2^8$	III	7	1
0	1	0	-2	-2	$+2^7$	II	7	3
0	1	0	$-3 + 2\epsilon$	$-3 + 2\epsilon$	$-\epsilon^{-5}2^8$	III	7	17
0	1	0	$-5 - 8\epsilon$	$3 + 8\epsilon$	$+\epsilon^5 2^{14}$	III^*	7	1
0	1	0	-9	7	$+2^{13}$	I_2^*	7	3
0	1	0	$-13 + 8\epsilon$	$11 - 8\epsilon$	$-\epsilon^{-5}2^{14}$	III^*	7	17
0	1	0	$-33 - 42\epsilon$	$103 + 158\epsilon$	$+\epsilon^{-1}2^8$	III	7	41
0	1	0	$-75 + 42\epsilon$	$261 - 158\epsilon$	$-\epsilon 2^8$	III	7	40
0	1	0	$-133 - 168\epsilon$	$-957 - 1432\epsilon$	$+\epsilon^{-1}2^{14}$	III^*	7	41
0	1	0	$-301 + 168\epsilon$	$-2389 + 1432\epsilon$	$-\epsilon 2^{14}$	III^*	7	40
0	$1 + \epsilon$	0	$-\epsilon$	$-1 - 2\epsilon$	$+\epsilon^5 2^8$	III	7	6
0	$1 + \epsilon$	0	$-2 + 3\epsilon$	1	$-\epsilon^{-5}2^8$	III	7	13
0	$1 + \epsilon$	0	$-2 - \epsilon$	1	$+\epsilon^{-1}2^8$	III	7	23
0	$1 + \epsilon$	0	$-2 - 2\epsilon$	$-4 - 6\epsilon$	$+\epsilon^4 2^7$	II	7	10
0	$1 + \epsilon$	0	$-2 - 4\epsilon$	$-6 - 10\epsilon$	$+\epsilon^5 2^7$	II	7	18
0	$1 + \epsilon$	0	$-2 - 7\epsilon$	-1	$+\epsilon^5 2^{14}$	III^*	7	6
0	$1 + \epsilon$	0	$-3 - \epsilon$	ϵ	$+\epsilon^{-4}2^7$	II	7	29
0	$1 + \epsilon$	0	$-4 + 3\epsilon$	$-5 + 2\epsilon$	$-\epsilon 2^8$	III	7	2
0	$1 + \epsilon$	0	$-5 + 3\epsilon$	$-6 + 3\epsilon$	$+\epsilon^4 2^7$	II	7	7
0	$1 + \epsilon$	0	$-5 + 4\epsilon$	$5 - 3\epsilon$	$+\epsilon^{-4}2^7$	II	7	24
0	$1 + \epsilon$	0	$-7 + 6\epsilon$	$9 - 5\epsilon$	$-\epsilon^{-5}2^7$	II	7	33
0	$1 + \epsilon$	0	$-10 + 9\epsilon$	$-9 + 8\epsilon$	$-\epsilon^{-5}2^{14}$	III^*	7	13
0	$1 + \epsilon$	0	$-10 - 7\epsilon$	$-25 - 24\epsilon$	$+\epsilon^{-1}2^{14}$	III^*	7	23
0	$1 + \epsilon$	0	$-10 - 11\epsilon$	$11 + 16\epsilon$	$+\epsilon^4 2^{13}$	I_2^*	7	10
0	$1 + \epsilon$	0	$-10 - 19\epsilon$	$19 + 32\epsilon$	$+\epsilon^5 2^{13}$	I_2^*	7	18
0	$1 + \epsilon$	0	$-14 - 7\epsilon$	$-21 - 36\epsilon$	$+\epsilon^{-4}2^{13}$	I_2^*	7	29
0	$1 + \epsilon$	0	$-18 + 9\epsilon$	$31 - 16\epsilon$	$-\epsilon 2^{14}$	III^*	7	2
0	$1 + \epsilon$	0	$-22 + 9\epsilon$	$35 - 28\epsilon$	$+\epsilon^4 2^{13}$	I_2^*	7	7
0	$1 + \epsilon$	0	$-22 + 13\epsilon$	$-49 + 28\epsilon$	$+\epsilon^{-4}2^{13}$	I_2^*	7	24
0	$1 + \epsilon$	0	$-30 + 21\epsilon$	$-81 + 52\epsilon$	$-\epsilon^{-5}2^{13}$	I_2^*	7	33

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$1 + \epsilon$	0	$-58 - 88\epsilon$	$-358 - 574\epsilon$	$+\epsilon^{-1}2^7$	<i>II</i>	7	43
0	$1 + \epsilon$	0	$-147 + 90\epsilon$	$785 - 485\epsilon$	$-\epsilon 2^7$	<i>II</i>	7	42
0	$1 + \epsilon$	0	$-234 - 355\epsilon$	$2275 + 3648\epsilon$	$+\epsilon^{-1}2^{13}$	I_2^*	7	43
0	$1 + \epsilon$	0	$-590 + 357\epsilon$	$-6513 + 4004\epsilon$	$-\epsilon 2^{13}$	I_2^*	7	42
0	$1 - \epsilon$	0	ϵ	-1	$-\epsilon 2^8$	<i>III</i>	7	13
0	$1 - \epsilon$	0	$-\epsilon$	1	$+\epsilon^{-1}2^8$	<i>III</i>	7	1
0	$1 - \epsilon$	0	-2	$-2 + 2\epsilon$	$+\epsilon^2 2^7$	<i>II</i>	7	24
0	$1 - \epsilon$	0	$-2 + 2\epsilon$	$-4 + 2\epsilon$	$-\epsilon 2^7$	<i>II</i>	7	33
0	$1 - \epsilon$	0	$-2 + 5\epsilon$	$1 + 2\epsilon$	$-\epsilon 2^{14}$	<i>III</i> *	7	13
0	$1 - \epsilon$	0	$-2 - 3\epsilon$	$-7 + 2\epsilon$	$+\epsilon^{-1}2^{14}$	<i>III</i> *	7	1
0	$1 - \epsilon$	0	$-4 - 7\epsilon$	$7 + 12\epsilon$	$+\epsilon^5 2^8$	<i>III</i>	7	23
0	$1 - \epsilon$	0	$-5 - 8\epsilon$	$9 + 15\epsilon$	$+\epsilon^2 2^7$	<i>II</i>	7	29
0	$1 - \epsilon$	0	$-10 + \epsilon$	$5 - 6\epsilon$	$+\epsilon^2 2^{13}$	I_2^*	7	24
0	$1 - \epsilon$	0	$-10 + 9\epsilon$	$13 - 6\epsilon$	$-\epsilon 2^{13}$	I_2^*	7	33
0	$1 - \epsilon$	0	$-18 - 27\epsilon$	$-47 - 78\epsilon$	$+\epsilon^5 2^{14}$	<i>III</i> *	7	23
0	$1 - \epsilon$	0	$-22 - 31\epsilon$	$-63 - 98\epsilon$	$+\epsilon^2 2^{13}$	I_2^*	7	29
0	$1 - \epsilon$	0	$-147 - 237\epsilon$	$-1270 - 2055\epsilon$	$+\epsilon^5 2^7$	<i>II</i>	7	43
0	$1 - \epsilon$	0	$-192 + 117\epsilon$	$1099 - 680\epsilon$	$-\epsilon^{-5} 2^8$	<i>III</i>	7	40
0	$1 - \epsilon$	0	$-590 - 947\epsilon$	$10517 + 17030\epsilon$	$+\epsilon^5 2^{13}$	I_2^*	7	43
0	$1 - \epsilon$	0	$-770 + 469\epsilon$	$-10031 + 6210\epsilon$	$-\epsilon^{-5} 2^{14}$	<i>III</i> *	7	40
0	-1	0	1	-1	-2^8	<i>III</i>	7	5
0	-1	0	3	5	-2^{14}	<i>III</i> *	7	5
0	-1	0	$-1 - 2\epsilon$	$1 + 2\epsilon$	$+\epsilon^5 2^8$	<i>III</i>	7	1
0	-1	0	-2	2	$+2^7$	<i>II</i>	7	3
0	-1	0	$-3 + 2\epsilon$	$3 - 2\epsilon$	$-\epsilon^{-5} 2^8$	<i>III</i>	7	17
0	-1	0	$-5 - 8\epsilon$	$-3 - 8\epsilon$	$+\epsilon^5 2^{14}$	<i>III</i> *	7	1
0	-1	0	-9	-7	$+2^{13}$	I_2^*	7	3
0	-1	0	$-13 + 8\epsilon$	$-11 + 8\epsilon$	$-\epsilon^{-5} 2^{14}$	<i>III</i> *	7	17
0	-1	0	$-33 - 42\epsilon$	$-103 - 158\epsilon$	$+\epsilon^{-1} 2^8$	<i>III</i>	7	41
0	-1	0	$-75 + 42\epsilon$	$-261 + 158\epsilon$	$-\epsilon 2^8$	<i>III</i>	7	40
0	-1	0	$-133 - 168\epsilon$	$957 + 1432\epsilon$	$+\epsilon^{-1} 2^{14}$	<i>III</i> *	7	41
0	-1	0	$-301 + 168\epsilon$	$2389 - 1432\epsilon$	$-\epsilon 2^{14}$	<i>III</i> *	7	40
0	$-1 + \epsilon$	0	ϵ	1	$-\epsilon 2^8$	<i>III</i>	7	13
0	$-1 + \epsilon$	0	$-\epsilon$	-1	$+\epsilon^{-1} 2^8$	<i>III</i>	7	1
0	$-1 + \epsilon$	0	-2	$2 - 2\epsilon$	$+\epsilon^2 2^7$	<i>II</i>	7	24
0	$-1 + \epsilon$	0	$-2 + 2\epsilon$	$4 - 2\epsilon$	$-\epsilon 2^7$	<i>II</i>	7	33
0	$-1 + \epsilon$	0	$-2 + 5\epsilon$	$-1 - 2\epsilon$	$-\epsilon 2^{14}$	<i>III</i> *	7	13
0	$-1 + \epsilon$	0	$-2 - 3\epsilon$	$7 - 2\epsilon$	$+\epsilon^{-1} 2^{14}$	<i>III</i> *	7	1
0	$-1 + \epsilon$	0	$-4 - 7\epsilon$	$-7 - 12\epsilon$	$+\epsilon^5 2^8$	<i>III</i>	7	23
0	$-1 + \epsilon$	0	$-5 - 8\epsilon$	$-9 - 15\epsilon$	$+\epsilon^2 2^7$	<i>II</i>	7	29
0	$-1 + \epsilon$	0	$-10 + \epsilon$	$-5 + 6\epsilon$	$+\epsilon^2 2^{13}$	I_2^*	7	24
0	$-1 + \epsilon$	0	$-10 + 9\epsilon$	$-13 + 6\epsilon$	$-\epsilon 2^{13}$	I_2^*	7	33
0	$-1 + \epsilon$	0	$-18 - 27\epsilon$	$47 + 78\epsilon$	$+\epsilon^5 2^{14}$	<i>III</i> *	7	23
0	$-1 + \epsilon$	0	$-22 - 31\epsilon$	$63 + 98\epsilon$	$+\epsilon^2 2^{13}$	I_2^*	7	29
0	$-1 + \epsilon$	0	$-147 - 237\epsilon$	$1270 + 2055\epsilon$	$+\epsilon^5 2^7$	<i>II</i>	7	43
0	$-1 + \epsilon$	0	$-192 + 117\epsilon$	$-1099 + 680\epsilon$	$-\epsilon^{-5} 2^8$	<i>III</i>	7	40
0	$-1 + \epsilon$	0	$-590 - 947\epsilon$	$-10517 - 17030\epsilon$	$+\epsilon^5 2^{13}$	I_2^*	7	43
0	$-1 + \epsilon$	0	$-770 + 469\epsilon$	$10031 - 6210\epsilon$	$-\epsilon^{-5} 2^{14}$	<i>III</i> *	7	40

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$-1 - \epsilon$	0	$-\epsilon$	$1 + 2\epsilon$	$+\epsilon^5 2^8$	<i>III</i>	7	6
0	$-1 - \epsilon$	0	$-2 + 3\epsilon$	-1	$-\epsilon^{-5} 2^8$	<i>III</i>	7	13
0	$-1 - \epsilon$	0	$-2 - \epsilon$	-1	$+\epsilon^{-1} 2^8$	<i>III</i>	7	23
0	$-1 - \epsilon$	0	$-2 - 2\epsilon$	$4 + 6\epsilon$	$+\epsilon^4 2^7$	<i>II</i>	7	10
0	$-1 - \epsilon$	0	$-2 - 4\epsilon$	$6 + 10\epsilon$	$+\epsilon^5 2^7$	<i>II</i>	7	18
0	$-1 - \epsilon$	0	$-2 - 7\epsilon$	1	$+\epsilon^5 2^{14}$	<i>III</i> *	7	6
0	$-1 - \epsilon$	0	$-3 - \epsilon$	$-\epsilon$	$+\epsilon^{-4} 2^7$	<i>II</i>	7	29
0	$-1 - \epsilon$	0	$-4 + 3\epsilon$	$5 - 2\epsilon$	$-\epsilon 2^8$	<i>III</i>	7	2
0	$-1 - \epsilon$	0	$-5 + 3\epsilon$	$6 - 3\epsilon$	$+\epsilon^4 2^7$	<i>II</i>	7	7
0	$-1 - \epsilon$	0	$-5 + 4\epsilon$	$-5 + 3\epsilon$	$+\epsilon^{-4} 2^7$	<i>II</i>	7	24
0	$-1 - \epsilon$	0	$-7 + 6\epsilon$	$-9 + 5\epsilon$	$-\epsilon^{-5} 2^7$	<i>II</i>	7	33
0	$-1 - \epsilon$	0	$-10 + 9\epsilon$	$9 - 8\epsilon$	$-\epsilon^{-5} 2^{14}$	<i>III</i> *	7	13
0	$-1 - \epsilon$	0	$-10 - 7\epsilon$	$25 + 24\epsilon$	$+\epsilon^{-1} 2^{14}$	<i>III</i> *	7	23
0	$-1 - \epsilon$	0	$-10 - 11\epsilon$	$-11 - 16\epsilon$	$+\epsilon^4 2^{13}$	I_2^*	7	10
0	$-1 - \epsilon$	0	$-10 - 19\epsilon$	$-19 - 32\epsilon$	$+\epsilon^5 2^{13}$	I_2^*	7	18
0	$-1 - \epsilon$	0	$-14 - 7\epsilon$	$21 + 36\epsilon$	$+\epsilon^{-4} 2^{13}$	I_2^*	7	29
0	$-1 - \epsilon$	0	$-18 + 9\epsilon$	$-31 + 16\epsilon$	$-\epsilon 2^{14}$	<i>III</i> *	7	2
0	$-1 - \epsilon$	0	$-22 + 9\epsilon$	$-35 + 28\epsilon$	$+\epsilon^4 2^{13}$	I_2^*	7	7
0	$-1 - \epsilon$	0	$-22 + 13\epsilon$	$49 - 28\epsilon$	$+\epsilon^{-4} 2^{13}$	I_2^*	7	24
0	$-1 - \epsilon$	0	$-30 + 21\epsilon$	$81 - 52\epsilon$	$-\epsilon^{-5} 2^{13}$	I_2^*	7	33
0	$-1 - \epsilon$	0	$-58 - 88\epsilon$	$358 + 574\epsilon$	$+\epsilon^{-1} 2^7$	<i>II</i>	7	43
0	$-1 - \epsilon$	0	$-147 + 90\epsilon$	$-785 + 485\epsilon$	$-\epsilon 2^7$	<i>II</i>	7	42
0	$-1 - \epsilon$	0	$-234 - 355\epsilon$	$-2275 - 3648\epsilon$	$+\epsilon^{-1} 2^{13}$	I_2^*	7	43
0	$-1 - \epsilon$	0	$-590 + 357\epsilon$	$6513 - 4004\epsilon$	$-\epsilon 2^{13}$	I_2^*	7	42
0	0	0	2ϵ	0	$-\epsilon^3 2^9$	<i>III</i>	8	11
0	0	0	8ϵ	0	$-\epsilon^3 2^{15}$	<i>III</i> *	8	11
0	0	0	-2ϵ	0	$+\epsilon^3 2^9$	<i>III</i>	8	11
0	0	0	-8ϵ	0	$+\epsilon^3 2^{15}$	<i>III</i> *	8	11
0	0	0	2	0	-2^9	<i>III</i>	8	11
0	0	0	$2 + 2\epsilon$	0	$-\epsilon^6 2^9$	<i>III</i>	8	11
0	0	0	$2 - 2\epsilon$	0	$+\epsilon^{-3} 2^9$	<i>III</i>	8	11
0	0	0	8	0	-2^{15}	<i>III</i> *	8	11
0	0	0	$8 + 8\epsilon$	0	$-\epsilon^6 2^{15}$	<i>III</i> *	8	11
0	0	0	$8 - 8\epsilon$	0	$+\epsilon^{-3} 2^{15}$	<i>III</i> *	8	11
0	0	0	-2	0	$+2^9$	<i>III</i>	8	11
0	0	0	$-2 + 2\epsilon$	0	$-\epsilon^{-3} 2^9$	<i>III</i>	8	11
0	0	0	$-2 - 2\epsilon$	0	$+\epsilon^6 2^9$	<i>III</i>	8	11
0	0	0	-8	0	$+2^{15}$	<i>III</i> *	8	11
0	0	0	$-8 + 8\epsilon$	0	$-\epsilon^{-3} 2^{15}$	<i>III</i> *	8	11
0	0	0	$-8 - 8\epsilon$	0	$+\epsilon^6 2^{15}$	<i>III</i> *	8	11
0	ϵ	0	$-3 - \epsilon$	-1	$+\epsilon^6 2^9$	<i>III</i>	8	26
0	ϵ	0	$-3 - 3\epsilon$	$1 + 2\epsilon$	$+\epsilon^6 2^9$	<i>III</i>	8	35
0	ϵ	0	$-3 - 5\epsilon$	$3 + 4\epsilon$	$+\epsilon^3 2^9$	<i>III</i>	8	16
0	ϵ	0	$-5 - 7\epsilon$	$5 + 8\epsilon$	$+\epsilon^6 2^9$	<i>III</i>	8	12
0	ϵ	0	$-13 - 5\epsilon$	$3 - 18\epsilon$	$+\epsilon^6 2^{15}$	<i>III</i> *	8	26
0	ϵ	0	$-13 - 13\epsilon$	$-21 - 42\epsilon$	$+\epsilon^6 2^{15}$	<i>III</i> *	8	35
0	ϵ	0	$-13 - 21\epsilon$	$-45 - 66\epsilon$	$+\epsilon^3 2^{15}$	<i>III</i> *	8	16
0	ϵ	0	$-21 - 29\epsilon$	$-69 - 114\epsilon$	$+\epsilon^6 2^{15}$	<i>III</i> *	8	12

a_1	a_2	a_3	a_4	a_6	Δ	type	f	code
0	$-\epsilon$	0	$-3 - \epsilon$	1	$+\epsilon^6 2^9$	III	8	26
0	$-\epsilon$	0	$-3 - 3\epsilon$	$-1 - 2\epsilon$	$+\epsilon^6 2^9$	III	8	35
0	$-\epsilon$	0	$-3 - 5\epsilon$	$-3 - 4\epsilon$	$+\epsilon^3 2^9$	III	8	16
0	$-\epsilon$	0	$-5 - 7\epsilon$	$-5 - 8\epsilon$	$+\epsilon^6 2^9$	III	8	12
0	$-\epsilon$	0	$-13 - 5\epsilon$	$-3 + 18\epsilon$	$+\epsilon^6 2^{15}$	III*	8	26
0	$-\epsilon$	0	$-13 - 13\epsilon$	$21 + 42\epsilon$	$+\epsilon^6 2^{15}$	III*	8	35
0	$-\epsilon$	0	$-13 - 21\epsilon$	$45 + 66\epsilon$	$+\epsilon^3 2^{15}$	III*	8	16
0	$-\epsilon$	0	$-21 - 29\epsilon$	$69 + 114\epsilon$	$+\epsilon^6 2^{15}$	III*	8	12
0	1	0	$-1 - 2\epsilon$	$-1 + 2\epsilon$	$+\epsilon^{-3} 2^9$	III	8	16
0	1	0	-3	1	$+2^9$	III	8	35
0	1	0	$-3 + 2\epsilon$	$1 - 2\epsilon$	$-\epsilon^3 2^9$	III	8	32
0	1	0	$-3 - 2\epsilon$	$1 + 2\epsilon$	$+2^9$	III	8	12
0	1	0	$-5 + 2\epsilon$	$3 - 2\epsilon$	$+2^9$	III	8	26
0	1	0	$-5 - 8\epsilon$	$3 - 24\epsilon$	$+\epsilon^{-3} 2^{15}$	III*	8	16
0	1	0	-13	-21	$+2^{15}$	III*	8	35
0	1	0	$-13 + 8\epsilon$	$-21 + 24\epsilon$	$-\epsilon^3 2^{15}$	III*	8	32
0	1	0	$-13 - 8\epsilon$	$-21 - 24\epsilon$	$+2^{15}$	III*	8	12
0	1	0	$-21 + 8\epsilon$	$-45 + 24\epsilon$	$+2^{15}$	III*	8	26
0	$1 - \epsilon$	0	$-8 + 5\epsilon$	$7 - 4\epsilon$	$-\epsilon^{-3} 2^9$	III	8	32
0	$1 - \epsilon$	0	$-34 + 21\epsilon$	$-111 + 66\epsilon$	$-\epsilon^{-3} 2^{15}$	III*	8	32
0	-1	0	$-1 - 2\epsilon$	$1 - 2\epsilon$	$+\epsilon^{-3} 2^9$	III	8	16
0	-1	0	-3	-1	$+2^9$	III	8	35
0	-1	0	$-3 + 2\epsilon$	$-1 + 2\epsilon$	$-\epsilon^3 2^9$	III	8	32
0	-1	0	$-3 - 2\epsilon$	$-1 - 2\epsilon$	$+2^9$	III	8	12
0	-1	0	$-5 + 2\epsilon$	$-3 + 2\epsilon$	$+2^9$	III	8	26
0	-1	0	$-5 - 8\epsilon$	$-3 + 24\epsilon$	$+\epsilon^{-3} 2^{15}$	III*	8	16
0	-1	0	-13	21	$+2^{15}$	III*	8	35
0	-1	0	$-13 + 8\epsilon$	$21 - 24\epsilon$	$-\epsilon^3 2^{15}$	III*	8	32
0	-1	0	$-13 - 8\epsilon$	$21 + 24\epsilon$	$+2^{15}$	III*	8	12
0	-1	0	$-21 + 8\epsilon$	$45 - 24\epsilon$	$+2^{15}$	III*	8	26
0	$-1 + \epsilon$	0	$-8 + 5\epsilon$	$-7 + 4\epsilon$	$-\epsilon^{-3} 2^9$	III	8	32
0	$-1 + \epsilon$	0	$-34 + 21\epsilon$	$111 - 66\epsilon$	$-\epsilon^{-3} 2^{15}$	III*	8	32

Table 2. Elliptic curves defined over $\mathbb{Q}(\sqrt{5})$ with good reduction away from 2.

REFERENCES

- [1] B.J. Birch and W. Kuyk (eds.), *Modular functions of one variable IV*, Lecture notes in Mathematics, vol. 476, Berlin, Springer-Verlag, 1975.
- [2] J.H.E. Cohn, *Lucas and Fibonacci numbers and some Diophantine equations*, Proc. Glasgow Math. Assoc. **7** (1965), 24–28.
- [3] J.E. Cremona, *Modular symbols for $\Gamma_1(N)$ and elliptic curves with everywhere good reduction*, Math. Proc. Cambridge Philos. Soc. **111** (1992), 199–218.
- [4] J.E. Cremona, *Algorithms for modular elliptic curves*, Cambridge University Press, Cambridge, 1992.
- [5] H. Ishii, *The non-existence of elliptic curves with everywhere good reduction*, J. Maths Soc. Japan **31** (1979), 273–279.
- [6] ———, *The non-existence of elliptic curves with everywhere good reduction over certain quadratic fields*, Japan. J. Maths. **12** (1986), 45–52.
- [7] R.G.E. Pinch, *Elliptic curves with everywhere good reduction*, Submitted.
- [8] ———, *Elliptic curves with good reduction away from 2*, Math. Proc. Cambridge Philos. Soc. **96** (1984), 25–38.
- [9] ———, *Elliptic curves with good reduction away from 2: II*, Math. Proc. Cambridge Philos. Soc. **100** (1986), 435–457.
- [10] ———, *Simultaneous Pellian equations*, Math. Proc. Cambridge Philos. Soc. **103** (1988), 35–46.
- [11] ———, *Squares in quadratic progression*, Math. Comp. **60** (1993), 841–845.
- [12] J. Tate, *Algorithm for determining the type of a singular fibre in an elliptic pencil*, In Birch and Kuyk [1], pp. 33–52.

QUEENS' COLLEGE, UNIVERSITY OF CAMBRIDGE, SILVER STREET, CAMBRIDGE CB3 9ET, U.K.
E-mail address: `rgep@cam.ac.uk`